IMPERIAL STEREOSCOPIC ANATOMY OF THE HEAD AND NECK

SIMPLE MASTOIDECTOMY


Section-X-No. 3

SCHWARTZE ZAUFAL

Operation for Acute Mastoiditis

This operation is performed after the mastoid antrum has been opened, as shown in stereogram No. 2. The entire external table is removed (never going above the superior limit line A of the square, see stereogram No. 2), including the mastoid tip, with complete ablation of all the mastoid cells (zygomatic, squamous and mastoid) eniarging the antral mouth of the aditus ad. antrum. Ossicles are not disturbed.

This specimen has been tilted slightly backward to show more distinctly the antral tegmen. The head of the malleus is just visible, anterior to the incus, and the short incal process as it lies in its groove on the aditus floor is easily seen.



The figures indicate-

1—Antral Tegmen
2—Mas.oid Antrum (Floor)
3—Eminence Ext. Semi-Cir. Canal
4—Incus
5—Tympanic Tegmen
6—Aditus Ad. Antrum (Floor)

7-Fosterior Wall of Auditory Canal

8-Tympanum

9-Frominence of Lateral Sinus

Copyrighted, 1911, by Imperial Publishing Co. IMPERIAL PUBLISHING COMPANY - - - - NEW YORK

