OWER LIMB-SCARPA'S TRIANGLE. No. 1. (EDINBURGH STEREOSCOPIC ANATOMY. Copyright: T. C. & E. C. Jack, Edinburgh, & 34 Henrietta St., London, W.C.

LOWER LIMB.

SCARPA'S TRIANGLE-No. 1.

THE SKIN, LAYERS OF FASCIA, CUTANEOUS NERVES AND VESSELS, ETC., WHICH FORM THE ROOF OF THE SPACE, HAVE BEEN REMOVED, TO SHOW THE BOUNDARIES AND CONTENTS OF THE TRIANGLE.

Scarpa's triangle is a triangular space situated in the upper part of the front of the thigh. It is bounded above by Poupart's ligament, externally by the sartorius, and internally by the

inner margin of the adductor longus muscles.

POUPART'S LIGAMENT is a strong fibrous band extending from the anterior superior spine of the ilium to the spine of the pubis. At the inner end some fibres pass backwards and outwards and

gain an attachment to the ilio-pectineal line. This portion is known as Gimbernat's ligament.

The SARTORIUS arises from the anterior superior spine of the ilium, and from the notch below it, and consists of parallel fleshy fibres which form a long, strap-like muscle, which passes obliquely across the front of the thigh, and the ADDUCTOR LONGUS takes origin by a rounded tendon from the front of the pubis, in the angle between the crest and symphisis, from which the fleshy fibres diverge in a fan-shaped manner, to an insertion into the inner lip of the linea aspera.

Within the space are found the femoral artery and vein and the anterior crural nerve, with branches derived from them, the termination of the long saphenous vein, the crural branch of the genito-crural nerve, and some deep femoral lymphatic glands, together with areolar and fatty tissue.

On the inner side of the femoral vein is the femoral or crural ring, through which a femoral hernia may occur. It is a narrow space bounded by Poupart's ligament in front, by the fascia of the pectineus muscle and the pubic bone behind, by Gimbernat's ligament internally, and by the femoral vein externally, from which it is separated by a thin septum. It is filled by some fatty tissue and a lymphatic gland.

The roof of Scarpa's triangle is seen in the views of the Inguinal Region (Section 2).

1. Femoral artery.

- 2. Femoral vein.
- 3. Bifurcation of the femoral artery, and branches of anterior crural nerve.

The figures indicate—

- 4. Deep external pudic vessels.
- 5. Cutaneous branches from anterior crural nerve.
- 6. Poupart's ligament.
- 7 Iliacus muscle

- 8. Pectineus muscle.
- 9. Sartorius muscle.
- 10. Adductor longus muscle.
- 11. Rectus femoris muscle.
- 12. Ilio-tibial band.

EDINBURGH: T. C. & E. C. JACK; AND 34 HENRIETTA STREET, LONDON, W.C.



