

Phacoides crenulatus Conrad.
Phacoides radians Conrad.
Phacoides trisulcatus Conrad.
Phacoides tuomeyi Dall.
Plicatula marginata Say.
Rangia clathrodonta Conrad.
Semele bella Conrad.
Sportella constricta Conrad.

Sportella protexta Conrad.
Strigella flexuosa Say.
Tagelus gibbus Spengler.
Transenella carolinensis Dall.
Venericardia granulata Say.
Venericardia perplana Conrad.
Venericardia tridentata Say.
Venus rileyi Conrad.

WALKER'S BLUFF, RIGHT BANK OF CAPE FEAR RIVER, 60 MILES ABOVE
 WILMINGTON.

Pleistocene:

	FEET. INCHES.	
Sandy loam	2	
Mottled reddish and yellowish arenaceous clay.....	1	6
Drab clay interstratified with yellowish to reddish sand....	2	
Laminated drab clay, becoming arenaceous and with fine sand partings in lower half; basal 12 inches iron stained.	11	

Pliocene. (Waccamaw) :

Coarse, loose, orange-colored sand.....	3	
Calcareous tough clay.....	...	6
Shell marl full of fossils, many perfectly preserved. The perfect specimens are in a matrix of fine shell fragments mixed in places with coarse buff sand. Thickens in places through dipping down in pockets or holes in Cretaceous. Gradually thins out and disappears in middle of bluff through being cut out by overlying Pleistocene. Again appears at bend. It contains some phosphate nodules and some sandstone cobbles 6 inches in diameter. The average thickness is about 5-6 feet, maximum.....	12	

Cretaceous:

Laminated clays and sands, containing Cretaceous fossils in places, exposed to water's edge.....	48-51
--	-------

FOSSILS FROM WALKER'S BLUFF.

GASTROPODA :

Conus marylandicus Green.
Crepidula aculeata Gmelin var.
costata Morton.
Crepidula fornicata Say.
Crepidula plana Say.
Eupleura caudata Say.
Fulgur canaliculatum Linné.
Fulgur perversum Linné.
Marginella limatula Conrad.
Nassa caloosaensis Dall.
Nassa lapenotierei Dall.
Nassa trivitatta Say.
Nassa vibex Say.
Oliva litterata Lamarck.
Polynices duplicatus Say.

Polynices heros Say.
Seila adamsi H. C. Lea.
Terebra concava Say.
Terebra dislocata Say.
Terebra protexta Conrad.
Tornatina canaliculata Say.
Turbonilla interrupta Totten.
Turbonilla reticulata C. B. Adams.

PELECYPODA :

Abra aequalis Say.
Aligena aequata Conrad.
Anomia simplex d'Orbigny.
Arca adamsi Smith.
Arca limula Conrad.
Arca plicatula Conrad.
Arca rustica Tuomey & Holmes.