

CALCULI, RENAL AND VESICAL, WITH ALKALINE URINE

(Continued).

- 138—R Chloroformi,
Ext. hyoscyami, āā ℥viiss.
Vini opii, ℥iiss.
Olei anthemidis, ℥ix ℥iij.—M.

Ft. linimentum.

Sig.: Apply over the whole abdomen, and rub in well. RELIQUET.

CANCER.

- 139—R Condurango corticis contus., ℥iiss.
Syrupi simplicis, ℥v.
Aq. bullientis, ad ℥vj.—M.

Ft. infusio.

Sig.: A tablespoonful every hour or two, the whole to be taken during the day. To be continued for several months. (*In gastric cancer.*)

L. RIESS, Annual Univ. Med. Sci.

- 140—R Antifebrin., ℥j.

In capsulas no. xii div.

Sig.: Take a capsule, and repeat in twenty minutes, if required. (*For lancinating pains and insomnia of cancer.*) DEMIÉVILLE, Annual Univ. Med. Sci.

- 141—R Ext. conii fructus fld., ℥ss.

Sig.: Take ten minims every half hour till sleep comes on. (*For pain and insomnia of cancer. The preparation must be good, and made from the unripe fruit without heat.*)

MADIGAN, Annual Univ. Med. Sci.

- 142—R Bismuthi subnitrat., ℥ij.
Morphiæ sulph., gr. j.—M.

In pulveres no. vi div.

Sig.: A powder three times daily in milk when gastric pain and vomiting. (*In gastric cancer.*)

BARTHOLOW.

- 143—R Zinci chloridi, ℥j.
Pulv. althææ radiceis, ℥iij.
Aquæ, q. s. ut ft. magna.—M.

Sig.: Apply locally. (*In epithelioma.*)

CANQUOIN.

- 144—R Alveloz, ℥j.

Sig.: Use as a topical dressing. (*In uterine and superficial carcinomata.*)

BARNSFATHER, Annual Univ. Med. Sci.