

CHANCROID (Continued).

199—R Succī limonis, ℥^{iss}.
Vini opii, ℥^{xlv}.
Liq. plumbi subacet., ℥^j.
Aquæ destillat., ℥^v.—M.

Ft. lotio.

Sig.: Soak pledgets of lint in the solution and apply locally. (*In phagedenic chaneroid.*) RODET.

CHILBLAINS.

200—R Acidi carbolicī, gr. x.
Cosmolini,
Olei terebinthinæ, āā ℥^j.—M.

Sig.: Apply to affected part.

DAVIDSON, Annual Univ. Med. Sci.

201—R Linimenti chloroformi, ℥^{ij}.

Sig.: Apply to part with gentle friction. (*Early stage.*) DAVIDSON, Annual Univ. Med. Sci.

202—R Tinct. iodiniī, ℥^j.

Apply to parts with brush. (*When swollen or granulated.*) DAVIDSON, Annual Univ. Med. Sci.

203—R Camphoræ, gr. lxxv.
Spts. vini rectific., ℥^{ijj}.
Glycerinæ, ℥^v.—M.

Ft. linimentum.

Use locally several times daily. (*When skin is not broken.*) Foy.

204—R Bismuthi salicylat., ℥^{ij}.
Pulv. amyli, ℥^{xviiij}.—M.

Sig.: First bathe the chilblains in a decoction of walnut-leaves, then rub with spirits of camphor and cover with the powder. To quiet the itching use the following:—

205—R Glycerinæ,
Aquæ rosæ, āā ℥^j.
Acidi tannici, gr. j.—M.

Sig.: Use as a lotion, and then dust on the above powder. E. BESNIER.