

GOITRE (Continued).

547—R Iodoformi, ʒj.
Adipis, ʒj.—M.

Ft. ungt.

Sig.: Apply locally.

WARING.

548—R Potassii bromidi, ʒss.

In pulv. no. xii div.

Sig.: A powder in a half-tumblerful of water three times daily. (*In exophthalmic goitre.*)

JON. HUTCHINSON.

549—R Tinct. iodinii, ʒj.

Sig.: Inject an hypodermic syringe-ful into the tumor every seven or eight days. After two or three weeks, fortnightly injections until cured. (*Continue from two to six months.*)

DUGUET, Annual Univ. Med. Sci.

GONORRHOEA.

550—R Potassii citratis, ʒss-j.
Spts. limonis, ʒss.
Syr. simplicis, ʒij.
Aquæ, ʒj.—M.

Sig.: A dessertspoonful, well diluted, three or four times daily, fasting. (*In first stage.*)

VAN BUREN and KEYES.

551—R Balsami copaibæ, ʒj.
Spts. ætheris nitrosi, āā ʒj.
Liq. potassæ, ʒij.
Ext. glycyrrhizæ, ʒss.
Misce et adde
Olei gaultheriæ, gtt. xvj.
Syr. acaciæ, ʒvj. —M.

Sig.: Tablespoonful three times daily. BUMSTEAD.

552—R Zinci sulphat., gr. i-ij.
Liq. plumbi subacet. dil., ʒj. —M.

Sig.: Shake, and inject three or four times daily, or—

553—R Liq. plumbi subacet. dil., ʒj.
Ext. opii aquos., gr. vj.—M.

Sig.: Use as injection two to four times daily.

VAN BUREN and KEYES.

554—R Potassii permanganatis, gr. i-ij.
Aquæ destillatæ, ʒj. —M.

Sig.: Use as injection. (*Gleet.*)

VAN BUREN and KEYES.