

MAMMARY INFLAMMATION (Continued).

830—R Morphiae, . . . gr. x.
Hydrargyri oleatis, . . . ℥ss.
Acidi oleici, . . . ℥ixss.—M.

Sig.: Apply thrice daily. Ten to thirty drops are sufficient.
JOHN MARSHALL.

831—R Gran. efferv. magnesi citratis
(Keasbey & Mattison), . . . ℥iv.

Sig.: A tablespoonful in water, followed by ten grains of quiniæ if there be fever. (*In incipient mammitis.*)
L. STARR, Annual Univ. Med. Sci.

832—R Cerati resinæ co., . . . ℥j.
Olei olivæ, . . . ℥i-ij.—M.

Ft. ungt.

Sig.: Apply, spread generously on soft linen or sheet-lint. (*When suppuration is threatened. Is as satisfactory as a poultice, and more easily managed.*)
WITHERSTINE.

MANIA, ACUTE.

833—R Methyal, . . . ℥ij.
Syr. aurantii cort., . . . ℥ij.
Aquæ, . . . ad ℥iv.—M.

Sig.: From a teaspoonful to a tablespoonful, repeated if necessary, to produce quietness or sleep.

MAIRET and COMBEMALE, Annual Univ. Med. Sci.

834—R Amyl hydratis, . . . ℥j.

Sig.: Forty to seventy-five minims in sweetened water, as required.

VON MERING, Annual Univ. Med. Sci.

835—R Ext. conii fld., . . .
Ext. hyoscyami fld., . . . āā ℥vij.
Chloral hydratis, . . . gr. x.
Aquæ, . . . ℥ij.—M.

Ft. haustus.

Sig.: To be taken at a draught, and repeated if required.
MADIGAN, Annual Univ. Med. Sci.

836—R Potassii bromidi, . . . ℥j.
Tinct. cannabis indicæ, . . . ℥j.
Syr. simplicis, . . . ℥ij.
Aquæ, . . . q. s. ad ft. ℥iv.—M.

Sig.: A tablespoonful thrice daily. (*In periodical mania and senile mania.*)
CLOUSTON.