

OTITIS, OTORRHŒA, AND OTALGIA (Continued).

963—R Acidi carbolici, 3j.
Glycerinæ, 3ix.—M.

Sig.: Instill a few drops into the ear two or three times daily, after cleansing. HARTMANN.

964—R Sol. boroglyceride (50 per cent.), 3j.

Sig.: Instill a few drops into the ear, after cleansing it, twice or thrice daily. L. W. FOX.

965—R Liquor. hydrogenii peroxidi (10-vol.), 3iv.

Sig.: Syringe the ear carefully with one part solution to two of water, and when cleansed instill a few drops of the above solution. C. H. BURNETT.

966—R Ungt. hydrarg. nitratis rubri, 3ss.

Sig.: Apply a small quantity to the affected skin. (*In chronic inflammation of external meatus.*) BARTHOLOW.

967—R Sodii carbonatis, ʒij.
Infusi picis liquidæ, Oj.—M.

Sig.: Inject and instill into the ear twice daily, using the following after each cleansing:—

968—R Picis liquidæ, ʒj.
Vaselini, ʒv.
Sulphuris loti, gr. vj.
Spts. camphoræ, gr. iv.
Chloral hydratis, gr. ij.—M.

Ft. ungt.

Sig.: Anoint the external ear and auditory canal, morning and evening, after cleansing. Then twice weekly use—

969—R Tinct. iodinii, ℥xxiv.
Glycerinæ,
Aquæ, āā 3j.
Potassii iodidi, gr. vj.
Vini opii, gr. xxiv.—M.

Sig.: Bathe the whole surface of the auditory canal and the external ear two or three times weekly. (*For eczematous inflammation of the external ear.*) MIOT.