

SKIN DISEASES (Continued).

1411—R Lactis recentis, . . . 3xiiss.
Glycerinæ, . . . 3viiss.
Acidi hydrochlorici, . . . ℥lxxv.
Ammonii muriatis, . . . 3j. —M.

Ft. lotio.

Sig.: Apply morning and evening with a camel's hair brush. (*In tan and freckles.*) MONIN.

1412—R Sodii hyposulphitis, . . . 3iij.
Acidi sulphurosi dil., . . . 3ss.
Aquæ, . . . q. s. ad Oj.—M.

Ft. lotio.

Sig.: Apply thoroughly to the scalp, to loosen the crusts. (*In tinea favosa.*) STARTIN.

1413—R Iodinii, . . . gr. x.
Potassii iodidi, . . . gr. xv.
Tinct. iodinii co., . . . 3j. —M.

Sig.: Apply to the scalp after the crusts have been removed by soaking in oil, poulticing, or by using the preceding lotion. (*In tinea favosa.*) TILBURY FOX.

1414—R Sulphuris loti, . . . 3j.
Olei cadini,
Hydrarg. chloridi corros., āā gr. v.—M.

Sig.: Apply four times daily, on the hairy portion of the skin and scalp. (*In tinea favosa.*) BAZIN.

1415—R Acidi salicylici,
Acidi chrysophanici, . . . āā 3ij gr. viiss.
Cretæ præparatæ, . . . 3ij gr. xlv.
Vaselini, . . . 3xviiss. —M.

Ft. ungt.

Sig.: Remove the crusts, epilate the hairs, and rub in the ointment for fifteen minutes at night. (*In tinea favosa.*) MONOE.

1416—R Iodinii, . . . 3i-ij.
Olei picis decoloratæ, . . . 3j. —M.

Ft. pasta.

Sig.: Apply every fourth or sixth day. When the mass begins to flake off, wash well, and re-apply the paste. (*In tinea tonsurans, or ringworm of the scalp. Also in tinea circinata.*) COSTER.