

Sw. Jerusalem 1889

Mrs. Wheeler # 5095

FORMULÆ.

ABORTION.

1—R Tinct. opii, ℥_{xx-xxx}.
Sig.: Mix with two or three tablespoonfuls of boiled starch and inject into the rectum. PARVIN.

2—R Mist. asafoetidæ, ℥_{viiij}.
Sig.: A tablespoonful several times daily. (*In habitual abortion.*)
NEGRI, Annual Univ. Med. Sci.

3—R Tinct. ferri chlor., ℥_{ss}.
Potassii chloratis, ℥_{iv}.
Syr. simplicis, ℥_j.
Aquæ menth. pip., ad ℥_{iv}.—M.
Sig.: A dessertspoonful in a wineglassful of water after meals. (*When due to fatty degeneration of the placenta.*)
STROTHER, Annual Univ. Med. Sci.

ABSCESSSES.

4—R Iodoformi, ℥_{iiss-3v}.
Ætheris, ℥_{vj}. —M.
Inject three to five ounces after aspirating the abscess. (*In cold or tubercular abscess.*)
MOSETIG-MOORHOF, Annual Univ. Med. Sci.

5—R Iodoformi, ℥_{ij}.
Glycerinæ, ℥_{iiss}.—M.
Inject the abscess cavity, after evacuating the pus. (*In cold or tubercular abscess.*)
BILLROTH.

6—R Calcii sulphidi, gr. ij.
Sacch. lactis, ℥_j.—M.
In chartulas no. xx div.
Sig.: Take one powder every hour or two.
RINGER.