

ACIDITY (See also Pyrosis).

7—℞ Sodii bicarb., ʒiij.
In pulveres no. xii dividenda.
A powder in a wineglassful of water after meals.
ALONZO CLARK.

8—℞ Sodii bicarb., ʒj.
Pulv. rhei, ʒss.
Spts. menth. pip., fʒij.
Aquæ, q. s. ad ʒiv.—M.
Sig.: A tablespoonful after meals. (*For acidity
combined with constipation.*) Bellevue Hosp.

9—℞ Pulv. ipecac., gr. ss.
Pulv. rhei, gr. ij.
Sodii bicarb., gr. xij.—M.
In chartulas no. xii dividenda.
Sig.: One powder every four to six hours to an
infant one year old. J. LEWIS SMITH.

10—℞ Tinct. nucis vomicæ, ʒj.
Sig.: Five drops a quarter of an hour before food
three times daily. RINGER.

11—℞ Infusi rhei, ʒiiss.
Sodii bicarb., ʒiiss.
Syr. aurantii cort., ʒvj.—M.
Sig.: A teaspoonful twice daily. EWALD.

ACNE (See also Skin Diseases).

12—℞ Liq. potassæ, ʒj.
Aquæ rosæ, ʒiv.—M.
Sig.: Apply with a soft sponge twice daily.
BARTHOLOW.

13—℞ Sulphuris iodidi, ʒss.
Adipis, ʒj.—M.
Sig.: Use freely over the eruption night and morn-
ing. (*In acne indurata and rosacea.*) RINGER.

14—℞ Naphthol β, ʒiiss.
Sulphur. precip., ʒss.
Vaselin. vel. lanolin.,
Sapon. viridis, āā ʒij.—M.
Leniter terendo fiat pasta.
Sig.: Spread a thin layer on affected skin and leave
for fifteen to twenty minutes; then rub off the oint-
ment and dust with powdered talc. LASSAR.