

ALBUMINURIA (Continued).

23—R Ol. erigerontis, ℥ss.

Sig.: Five drops on a lump of sugar every three or four hours. (*In the chronic forms.*)

BARTHOLOW, Annual Univ. Med. Sci.

24—R Auri et sodii chlor., gr. iij.
Hydrarg. chlor. corr., gr. v.
Ext. gentian., q. s.—M.

Ft. massa et in pil. no. lx div.

Sig.: One pill morning and evening.

BARTHOLOW.

25—R Sodii iodidi, gr. xv.
Sodii phosphat., ℥ss.
Sodii chloridi, ℥iij.
Aquæ, q. s. ad ft. sol.—M.

Sig.: To be taken in the course of the twenty-four hours.

SEMMOLA.

ALCOHOLISM.

26—R Sol. nitro-glycerin (1 per cent.), ℥ij.

Sig.: One drop every two hours. (*In acute form, with cerebral anæmia and intense depression.*)

VAN GOIDTSNOVEN, Annual Univ. Med. Sci.

27—R Strychniæ sulph., gr. j.
Aquæ font., ℥j.—M.

Sig.: Five minims, increased cautiously to twenty minims, hypodermically twice daily. (*In both acute and chronic forms.*)

DOBROBRAVOFF, Annual Univ. Med. Sci.

28—R Tinct. gentianæ co., ℥ij.
Tinct. calumbæ co., āā ℥ij.
Tinct. nucis vom., ℥lxxx.—M.

Sig.: A dessertspoonful before each meal.

LOOMIS.

29—R Spts. ammon. aromat., ℥ij.
Tinct. camphoræ, ℥iiss.
Tinct. hyoscyami, ℥iiss.
Spts. lavandulæ co., q. s. ad ℥ij.—M.

Sig.: A teaspoonful every hour till relieved. Then give—