

**ANGINA PECTORIS (Continued).**

59—R Pyridin., . . . . . ℥ss.

Sig.: Six to ten drops daily, increasing to twenty-five drops, well diluted with water. Or three to five drops may be inhaled.

DE RENZI, Annual Univ. Med. Sci.

60—R Zinci cyanidi, . . . . . gr. iv-v.  
Confect. rosæ, . . . . . q. s. —M.

In pilulas no. xl div.

Sig.: One pill three times daily.

LASHKEVITCH, Annual Univ. Med. Sci.

61—R Amyl nitritis, . . . . . ℥ij.

Sig.: Two to ten drops on handkerchief. For inhalation.

BRUNTON.

62—R Antipyrin., . . . . . ℥j.  
Syr. toltutan., . . . . . ℥j.  
Aquæ, . . . . . ad ℥ij.—M.

Sig.: A tablespoonful at intervals of one to four hours until relieved.

GERMAIN SÉE, Annual Univ. Med. Sci.

**ANTHRAX.**

63—R Acidi carbolici, . . . . . ℥<sub>xx-xxv</sub>.  
Aquæ, . . . . . ℥j. —M.

Inject a few drops into and around the pustule.

USPENSKI, Annual Univ. Med. Sci.

64—R Quiniæ, . . . . . ℥j.  
Ol. terebinth., . . . . . q. s. ut ft. pasta.—M.

Sig.: Apply to pustule.

F. RIVAS, Annual Univ. Med. Sci.

**APHTHÆ.**

65—R Potassii iodidi, . . . . . gr. i-v.  
Aquæ, . . . . . ℥j. —M.

Sig.: Apply locally.

BARTHOLOW.

66—R Mel. boracis, . . . . . ℥j.

Apply to patches with a brush.

RINGER.

67—R Potassii chloratis, . . . . . gr. x.  
Aquæ, . . . . . ℥j. —M.

Apply locally several times daily.

BRUNTON.