

BLADDER, AFFECTIONS OF (*See Catarrh*).

BOILS (*See Abscess*).

BRIGHT'S DISEASE (*See Albuminuria*).

BROMIDROSIS.

91—R Sodii biborat., ʒj.
Lanolini, ʒx.—M.

Sig.: For external use.
WULFSBERG, *Annual Univ. Med. Sci.*

92—R Ext. geranii mac. fld., . . . ʒij.

Sig.: For external use.
PEPPER, *Annual Univ. Med. Sci.*

BRONCHITIS.

93—R Vini ipecac., ʒij.
Vini antimonialis, ʒj.
Vini xerici, ʒij.—M.

Sig.: Three drops every hour to a child six months
old. (*Where larger tubes only are affected.*)
DESSAU, *Annual Univ. Med. Sci.*

94—R Tinct. aconiti rad., ʒss.

Sig.: One or two drops every hour. (*In severe
cases with fever, where medium and small tubes are
affected.*)
DESSAU, *Annual Univ. Med. Sci.*

95—R Terebene, ʒss.

Sig.: Two to five drops on sugar every four hours,
according to child's age. (*In chronic form.*)
CARMICHAEL, *Annual Univ. Med. Sci.*

96—R Terpene hydrate, ʒss.

Sig.: Two to five drops on sugar every four hours,
according to child's age. (*In chronic form. Is taste-
less and similar in effect to Terebene.*)
CAMMANN, *Annual Univ. Med. Sci.*

97—R Sol. hydrogen dioxidi (10-vol.), ʒij.

Sig.: A teaspoonful in a glassful of water three times
daily. (*In chronic bronchitis with dyspnœa.*)
DE BLEYER, *Annual Univ. Med. Sci.*