BRONCHITIS (Continued).

105—℞ Acidi hydrocyanici dil., . . ♏xvj.
Syr. pruni virginian.,
Aq. camphoræ, . . āā ℥j.—M.

Sig.: A teaspoonful every two or three hours. *(In violent, troublesome cough.)* HARTSHORNE.

106—℞ Ammonii carb., . . . ℈ij.
Spts. chloroformi, . . . ℥ss.
Inf. senegæ, ... ℥viij.—M.

Sig.: Two tablespoonfuls every four or six hours. FOTHERGILL.

107—℞ Tinct. sanguinariæ, . . . ʒj.
Tinct. lobeliæ, ... ʒj.
Vini ipecac., ... ʒij.
Syr. tolutan., ... ℥ss.—M.

Sig.: A teaspoonful every three hours. BARTHOLOW.

BUBO.

108—℞ Tinct. iodinii, ... ℥j.

Sig.: Apply with brush every other day till skin becomes tender. VAN BUREN.

109—℞ Sol. hydrogen peroxidi (10-vol.), ... ℥viij.

Sig.: Apply after suppuration has begun. RINGER.

110—℞ Cerati resinæ co., . . . ℥j.
Ol. olivæ, ... ʒi–ij.—M.

Sig.: Spread on lint and apply. *(To hasten suppuration and granulation.)* WITHERSTINE.

BUNIONS.

111—℞ Tinct. iodinii,
Tinct. belladonnæ, . . āā ʒij.—M.

Sig.: Apply twice daily with a brush. *Annual Univ. Med. Sci.*

112—℞ Acidi tannici,
Cosmolini, . . . āā ℥ss.—M.

Sig.: Apply to joint after the skin has been removed by blister. GROSS, *Annual Univ. Med. Sci.*