

CATARRH, GENITO-URINARY (Continued).

183—R Copaibæ,
 Spts. lavand. co., āā ℥ij.
 Mucil. acaciæ, ℥ss.
 Syrupi simp., ℥iij.
 Aquæ, ℥iv.—M.
 Sig.: A tablespoonful twice daily. WOOD.

184—R Infusi buchu, ℥vij.
 Potassii bicarb., ℥j.
 Tinct. hyoscyami, ℥iiss.
 Ext. sarsæ fl., ℥iv.—M.
 Sig.: Two tablespoonfuls three times daily. (*In irritable bladder, with acid urine.*) COULSON.

185—R Atropinæ sulph., gr. j.
 Acidi acetici, gtt. xx.
 Alcoholis,
 Aquæ, āā ℥ss.—M.
 Sig.: Four drops in a wineglassful of water before each meal. (*In acute cystitis.*) GOODELL.

186—R Ext. tritici fld.,
 Syr. amygdalæ, āā ℥ij.—M.
 Sig.: A dessertspoonful, in water, five or six times daily. (*In chronic cystitis.*) THOMPSON.

187—R Argenti nitrat., gr. vij.
 Aq. destillat., ℥iiiss.—M.
 Sig.: Inject into the bladder, every third or fourth day, after washing it out with warm water. RICORD.

188—R Iodoformi, gr. xij.
 Ext. hyoscyami, gr. viij.
 Olei theobromæ, ℥vj. —M.
 In suppositoria no. viii div.
 Sig.: Introduce one into the rectum twice daily, one hour after giving the patient a lukewarm-water enema. If there be an urethral discharge, take a terpine pill, gr. iss, night and morning. (*In gonorrhœal cystitis.*) RELIQUET.

189—R Fol. hyoscyami, ℥ss.
 Aq. bullientis, Oj.—M.
 Ft. infusio.
 Sig.: A tablespoonful every half-hour for one forenoon, unless throat becomes dry or patient drowsy. DIDAY.