

**CHLORODYNE.**

206—R Chloroformi purif., . . . . f ℥iv.  
Ætheris fort., . . . . f ℥j.  
Alcoholis fort.,  
Syrupi fusci, . . . . āā f ℥iv.  
Ext. glycyrrh. pulv., . . . . ℥iiss.  
Morph. muriat., . . . . gr. viij.  
Ol. menthæ pip., . . . . ℥xvj.  
Acidi hydrocyanici dil. (2 per  
cent.), . . . . f ℥ij.  
Syrupi, . . . . f ℥xviiss.—M.

Sig.: Ten to twenty drops as required. Always  
shake before using. P. SQUIRE.

**CHLOROSIS (See Anæmia).**

**CHOLERA.**

207—R Tinct. opii,  
Tinct. capsici,  
Tinct. rhei arom.,  
Spts. menthæ pip.,  
Spts. camphoræ, . . . . āā p. æq.—M.

Sig.: Twenty to forty minims, diluted.

RUSCHENBERGER.

208—R Acidi nitrosi, . . . . ℥j.  
Tinct. opii, . . . . gtt. xl.  
Aq. camphoræ, . . . . ℥viiij.—M.

Sig.: One-fourth part to be taken every three or  
four hours. HOPE.

209—R Strychniæ sulph., . . . . gr. ¼.  
Acidi sulphurici dil., . . . . ℥ss.  
Morphiæ sulph., . . . . gr. ij.  
Aq. camphoræ, . . . q. s. ad ℥iv.—M.

Sig.: A teaspoonful every hour or two, well diluted.  
(*In threatened collapse. Also as a prophylactic, given  
less frequently.*) BARTHOLOW.

210—R Tinct. opii,  
Tinct. capsici,  
Spts. camphoræ, . . . . āā ℥j.  
Chloroformi, . . . . ℥ij.  
Alcoholis, . . . q. s. ad ft. ℥v.—M.

Sig.: Twenty to forty minims, diluted. SQUIBB.

211—R Morphiæ sulph., . . . . gr. ij.  
Spts. camphoræ, . . . . ℥j. —M.

Sig.: Fifteen minims every three or four hours by  
mouth, or hypodermically, in severe cases. (*In initial  
stage.*) Also—