

CHOLERA (Continued).

212—R Morphiae sulph., gr. x.
Atropiae sulph., gr. j.
Aquae destillat., ℥x.—M.

Sig.: Five to ten minims hypodermically, as required. (*In stage of collapse.*)

NAKAMURA, *Annual Univ. Med. Sci.*

213—R Tinct. opii, ℥x-xv.
Chloroformi, ℥xv-xx.
Spts. vini rectific., ℥j.
Aquae, ℥j.—M.

Sig.: To be taken at one dose, and repeated every two or three hours until diarrhoea is checked.

LOWNDES.

CHOLERA INFANTUM.

214—R Tinct. cocæ (1 to 5), ℥j.

Sig.: Four to six drops every two hours at three months of age. Fifteen to twenty drops in older children.

DIEDERICHS, *Annual Univ. Med. Sci.*

215—R Naphthalini, gr. xx-lxx.
Ol. bergamii, gtt. i-ij. —M.

In pulv. no. xii div.

Sig.: A powder every two or three hours.

HOLT, *Annual Univ. Med. Sci.*

216—R Resorcin., gr. viii-xl.
Syr. aurantii cort., ℥j.
Aquae aurantii flor., ad ℥ij.—M.

Sig.: A teaspoonful every two hours.

FLIESBURG, *Annual Univ. Med. Sci.*

217—R Saloli, gr. vj.
Sacch. lactis, gr. x.—M.

In pulv. no. xii div.

Sig.: A powder every two hours. (*For a child aged six months. Between five and ten years old, five grains of salol may be taken every two hours.*)

GOELET, *Annual Univ. Med. Sci.*

218—R Hydrarg. chlorid. mit., gr. iij.
Cretæ præp., gr. xxxvj.
Plumbi acetat., gr. xij.
Pulv. ipecac., gr. iij.—M.

In chart. no. xii div.

Sig.: One every three hours.

CONDIE.