

CHOLERA INFANTUM (Continued).

219—R Tinct. opii deodorat., . . . gtt. xvj.
 Spts. ammon. aromat., . . . ℥j.
 Bismuth. subnitrat., . . . ℥ij.
 Syr. simplicis, . . . ℥iv.
 Mist. cretæ, . . . ℥iss.—M.

Sig.: Shake well, and give a teaspoonful every two or three hours to a child eight to twelve months old. (*Six months old, half the dose.*)
J. LEWIS SMITH.

220—R Hydrarg. chlorid. mit., . . . gr. ij.
 Sodii bicarb., . . . ℥j.
 Pulv. zingiberis, . . . gr. xij.—M.

In pulveres no. xii dividenda.

Sig.: One powder three or four times daily. (*In incipient stage.*)
HARTSHORNE.

221—R Potassii bromidi, . . . ℥ij.
 Syr. simplicis, . . . ℥ss.
 Aq. menth. pip., . . . ℥iss.—M.

Sig.: A teaspoonful every hour or two. (*When cerebral congestion or reflex irritation is the cause.*)
BARTHOLOW.

222—R Ol. ricini, . . . f℥ij.
 Pulv. acaciæ,
 Sacch. albi, . . . āā ℥ij.
 Tinct. opii, . . . ℥xxj.
 Aq. cinnam., . . . q. s. ad f℥iv.—M.

Sig.: A teaspoonful every two or three hours.
WEST.

223—R Hydrarg. cum cretæ, . . . gr. ij.
 Sacch. lactis, . . . gr. x.—M.

In pulv. no. xii div.

Sig.: A powder every hour.
RINGER.

CHORDEE.

224—R Ext. opii aquos., . . . gr. ij.
 Pulv. camphoræ, . . . gr. iv.—M.

In pil. no. ii div.

Sig.: One or both on retiring.
VAN BUREN and KEYES.

225—R Ext. opii aquos., . . . gr. iss.
 Ol. theobromæ, . . . gr. xxx.—M.

Ft. suppositor. no. i.

Sig.: Introduce into rectum on retiring.
VAN BUREN and KEYES.