

DIABETES INSIPIDUS (Continued).

13—R Pulv. opii, gr. iv.
Acidi gallici, ℥ij. —M.

In chart. no. xii div.

Sig.: One three or four times daily.

H. C. WOOD, JR.

314—R Codeinæ, gr. viij.
Syrupi,
Aquæ, āā ℥j.—M.

Sig.: A half-teaspoonful thrice daily, gradually increased to two teaspoonfuls.

PAVY.

DIABETES MELLITUS.

315—R Lithii carbonat., gr. xxx.
Sodii arseniat., gr. j.
Ext. gentianæ, gr. xv.—M.

Ft. massa et in pil. no. xx div.

Sig.: One pill morning and evening.

VIGIER, Annual Univ. Med. Sci.

316—R Potassii phosphat., gr. xvj.
Aquæ, ℥x. —M.

Sig.: A teaspoonful in a little wine or hot tea three times daily.

DUCHENNE, Annual Univ. Med. Sci.

317—R Morphiæ acetat., gr. viij.
Aquæ, ℥iv. —M.

Sig.: A teaspoonful four times daily, increasing rapidly, until in three months seven grains daily are taken. (To be used with restricted diet.)

BRUCE, Annual Univ. Med. Sci.

318—R Ext. belladonnæ, gr. xxx.
Ext. opii, gr. xv.—M.

Ft. massa et in pil. no. xx div.

Sig.: One three times daily, gradually increased to double the quantity. (To be used with restricted diet.)

VILLEMEN, Annual Univ. Med. Sci.

319—R Pepsinæ cryst., ℥j.

In pulv. no. xii div.

Sig.: One three times daily, after meals. Gradually increased, if necessary. (Restricted diet at first.)

GARDNER, Annual Univ. Med. Sci.