

DIABETES MELLITUS (Continued).

320—R Sodii salicylat., ℥iv-vj.
Glycerinæ, ℥j.
Aquæ, ad ℥iij.—M.

Sig.: Two teaspoonfuls three times daily.
DA COSTA, Annual Univ. Med. Sci.

321—R Sodii salicylat., ℥iij.
Liq. potassii arsenitis, ℥j.
Glycerinæ, ℥j.
Aquæ cinnamomi, ad ℥iij.—M.

Sig.: A dessertspoonful three times daily.
J. C. WILSON.

322—R Pulv. jambul. sem., ℥j.

Dispensa in capsulas no. xxiv.
Sig.: One or two capsules thrice daily, after food.
FENWICK, Annual Univ. Med. Sci.

323—R Sol. cocaini mur. (4 per cent.), ℥ss.

Sig.: Two drops every three hours. (*With anti-diabetic diet.*)

When polydipsia disappears, take with above—

324—R Tinct. opii, ℥j.
Tinct. ferri muriat., ℥ix.—M.

Sig.: Twenty drops three times daily.
WELLER, Annual Univ. Med. Sci.

325—R Iodoformi, gr. ij.

In pil. no. xii div.
Sig.: One pill after meals, three times daily. (*With restricted diet.*)
LEVI, Annual Univ. Med. Sci.

326—R Ergotini, ℥j.
Glycerinæ, ℥j.
Aquæ destillat., ℥vij.—M.

Sig.: Five or six drops daily, hypodermically. (*To diminish thirst.*)
CORNILLON, Annual Univ. Med. Sci.

DIARRHŒA, CHILDREN.

327—R Pulv. ipecac., gr. ss.
Pulv. rhei, gr. ij.
Sodii bicarb., gr. xij.—M.

In pulv. no. xii div.
Sig.: One powder every four to six hours to an infant one year old. (*In indigestion with acidity.*)
J. LEWIS SMITH.