

**DIARRHŒA, ADULTS (Continued).**

343—℞ Pil. hydrargyri, . . . . . gr. x.  
Morphiæ sulphat., . . . . . gr. j.  
Acidi tannici, . . . . . ℥j.  
Mucil. acaciæ, . . . q. s. ut ft. massa.—M.

Ft. massa et in pil. no. x div.

Sig.: A pill every two hours.

ELLIS.

344—℞ Saloli, . . . . . ℥ij.

In pulv. no. xii div.

Sig.: A powder every two hours, followed by a draught of water.

GOELET, Annual Univ. Med. Sci.

**DIPHThERIA.**

345—℞ Pilocarpinæ muriat., . . . . . gr.  $\frac{1}{3}$ — $\frac{3}{5}$ .  
Pepsinæ cryst., . . . . . gr. x—xij.  
Acidi hydrochlorici, . . . . . gtt. ij.  
Aquæ destillat., . . . . . ℥ij ℥iss.—M.

Sig.: A teaspoonful every hour, with a small amount of wine following it. (*For children.*)

LAX, Annual Univ. Med. Sci.

346—℞ Pilocarpinæ muriat., . . . . . gr. iss.  
Pepsinæ cryst., . . . . . gr. xxx.  
Acidi hydrochlorici, . . . . . gtt. iij.  
Aquæ destillat., . . . . . ℥viiij.—M.

Sig.: A tablespoonful every half hour in wine. (*For adults.*)

GUTTMAN, Annual Univ. Med. Sci.

347—℞ Hydrarg. cyanidi, . . . . . gr.  $\frac{3}{10}$ .  
Tinct. aconiti rad., . . . . . ℥xv.  
Aquæ destillat., . . . . . ℥xv.—M.

Sig.: One teaspoonful hourly.

LE GENDRE, Annual Univ. Med. Sci.

348—℞ Salinaphthol., . . . . . gr. xx—xxv.  
Spts. vini rectific., . . . . . ℥j. —M.

Sig.: A dessertspoonful in two-thirds of a tumbler of water as a gargle. Salinaphthol may also be used internally in doses of one to two drachms daily in children.

GEORGI, Annual Univ. Med. Sci.

349—℞ Tinct. iodinii, . . . . . gtt. viij.  
Potassii iodidi, . . . . . gr. iss.  
Syr. simplicis,  
Aquæ, . . . . . āā ℥ij.—M.

Sig.: To be taken in one dose. (*Prevention for exposed children.*)

DUMAS, Annual Univ. Med. Sci.