

DIPHTHERIA (Continued).

350—R Olei terebinthinæ, ℥iv.

Sig.: Put in a cup and set it in a basin of water on the top of the stove, so that the vapor fills the room. Renew when necessary. (*Prevention for exposed children.*)

LLEWELLYN ELLIOTT, Annual Univ. Med. Sci.

351—R Tinct. ferri chloridi, ℥ss.

Sig.: One drop every quarter of an hour. (*In phlegmonous form.*)

LUNIN, Annual Univ. Med. Sci.

352—R Olei terebinthinæ, ℥iv.

Sig.: A tablespoonful twice daily, or, in addition, ten drops every hour. (*In fibrinous forms.*)

LUNIN, Annual Univ. Med. Sci.

353—R Coal-tar, ℥vij.
Olei terebinthinæ, ℥ij ℥vj.—M.

Sig.: Light and keep burning in sick-room.

DELTHIL, Annual Univ. Med. Sci.

354—R Olei terebinthinæ, ℥ij.

Sig.: Ten minims to one teaspoonful, one to three times daily, in milk, sugar-water, or gruel, with alcoholic stimulants.

SCHENKER, Annual Univ. Med. Sci.

355—R Olei eucalypti, ℥ij.
Olei terebinthinæ, ℥viii.—M.

Sig.: Place in shallow vessels and keep boiling or at least simmering over the stove.

J. LEWIS SMITH, Annual Univ. Med. Sci.

356—R Tinct. ferri chloridi, ℥j.
Syr. simplicis, ℥ij.—M.

Sig.: A teaspoonful every hour to a child ten years old, or half a teaspoonful every half-hour.

FERGUSON, Annual Univ. Med. Sci.

357—R Tinct. ferri chlor., ℥j.
Glycerinæ,
Aquæ, āā ℥j.—M.

Sig.: A teaspoonful every hour.

BILLINGTON, Annual Univ. Med. Sci.