

DIPHTHERIA (Continued).

358—R Hydrarg. bichloridi, . . . gr. ss.
Spts. frumenti, . . . ʒj.
Syr. simplicis, . . . ʒj.—M.

Sig.: A teaspoonful every three hours, night and day.
DRESCHER, Annual Univ. Med. Sci.

359—R Hydrarg. chlor. cor., . . . gr. j.
Spts. vini rectific., . . . ʒij.
Elix. bismuthi et pepsinæ, ad ʒiv.—M.

Sig.: A teaspoonful every two hours for child six years old.

J. LEWIS SMITH, Annual Univ. Med. Sci.

360—R Acidi carbolici, . . . ʒx.
Acidi salicylici, . . . ʒij.
Acidi benzoici, . . . ʒiv.
Spts. vini rectificat., q. s. ad ft. sol.—M.

Sig.: To water constantly boiling add a spoonful of above solution, using the whole quantity above in twenty-four hours. The quantity may be increased if the size of the room, age of the patient, or severity of the disease require it.

RENOU, Annual Univ. Med. Sci.

361—R Hydrarg. chlorid. mit., . . . ʒj.

In pulv. no. xxiv div.

Sig.: One or two powders every one, two, or three hours until free catharsis follows, and then at longer intervals, so that three or four evacuations are produced daily.

W. C. RESTER, Annual Univ. Med. Sci.

362—R Tinct. ferri chloridi, . . . ʒii-ij.
Potassii chlorat., . . . ʒj.
Acidi muriatici dil., . . . gtt. x.
Syr. simplicis, . . . ʒiv.—M.

Sig.: A teaspoonful every hour or two.

J. LEWIS SMITH, Annual Univ. Med. Sci.

363—R Liq. ferri subsulph., . . . ʒij.
Glycerinæ, . . . ʒij.—M.

Sig.: Apply locally with brush twice daily.

DRESCHER, Annual Univ. Med. Sci.

364—R Acidi carbolici, . . . gtt. x.
Liq. ferri subsulph., . . . ʒij.
Glycerinæ, . . . ʒj.—M.

Sig.: To be applied every three to six hours with a camel's hair brush.

J. LEWIS SMITH, Annual Univ. Med. Sci.