

**DYSENTERY (Continued).**

**388—R** Thymoli, . . . . . ʒj.

Ft. massa et in pil. no. xxiv div.

Sig.: Two pills every six hours.

MARTINI, *Annual Univ. Med. Sci.*

**389—R** Plumbi acetat., . . . . . gr. xxiv.  
Pulv. ipecac., . . . . . gr. iij.  
Pulv. opii, . . . . . gr. iij.—M.

Ft. massa et in pil. no. xii div.

Sig.: One pill every two hours until blood ceases;  
then at longer intervals. DA COSTA.

**390—R** Naphthalini, . . . . . ʒiss.

Dispensa in capsulas no. xviiij.

Sig.: Two capsules every three hours. At least  
twelve to be taken during the twenty-four hours.

HOLT, *Annual Univ. Med. Sci.*

**391—R** Tinct. opii deodorat., . . . . . ʒss.  
Bismuthi subnitrat., . . . . . ʒij.  
Aquæ menthæ pip.,  
Syr. zingiberis, . . . . . āā ʒj.—M.

Sig.: Shake bottle. Give one teaspoonful every two  
to four hours, to a child five years of age. Half dose  
for child one year of age. J. LEWIS SMITH.

**392—R** Vini ipecac., . . . . . ʒss.

Sig.: One drop every hour. (*In acute or chronic  
form of children, with slimy stools.*) RINGER.

**DYSMENORRHŒA.**

**393—R** Potassii bromidi,  
Chloral hydrat., . . . . . āā ʒiv.  
Syr. simplicis,  
Aquæ, . . . . . āā ʒij.—M.

Sig.: Two tablespoonfuls to be used as an enema, as  
required, for pain. MENIÈRE, Annual Univ. Med. Sci.

**394—R** Antipyrin., . . . . . ʒij.  
Syr. toltutan., . . . . . ʒij.—M.

Sig.: Two teaspoonfuls at first; one teaspoonful  
every hour or two afterward until pain is relieved.

DELLENBAUGH, Annual Univ. Med. Sci.