

**DYSMENORRHŒA (Continued).**

395—R Antipyrin, . . . . . gr. xxviiiiss.  
Cocaini muriat., . . . . . gr. iss.  
Aquæ bullientis, . . . . . ℥ij. —M.

Sig.: Ten to twenty minims hypodermically. (*In neuralgic and congestive forms.*) May also be taken by mouth. P. MENIÈRE, Annual Univ. Med. Sci.

396—R Tinct. guaiaci ammoniat., . . . . . ℥ij.

Sig.: One-half to one teaspoonful in milk, every two or three hours, until pain is relieved.

SIR JAMES SAWYER, Annual Univ. Med. Sci.

397—R Tinct. pulsatillæ rad., . . . . . ℥ss.

Sig.: Two or three drops every two hours for ten days preceding the period.

BROWN, Annual Univ. Med. Sci.

398—R Ext. opii, . . . . . gr. v.  
Ext. cannabis ind., . . . . .  
Ext. hyoscyami, . . . . . āā gr. x.  
Pulv. camphoræ, . . . . . gr. xxv.—M.

Ft. massa et in pil. no. x div.

Sig.: A pill two or three times daily. McLANE.

399—R Liq. ammonii acetat., . . . . . ℥iv.

Sig.: A tablespoonful every two or three hours, with the following:—

400—R Pulv. ipecac., . . . . . gr. iv.

In pil. no. xii div.

Sig.: One every two or three hours. EMMET.

**DYSPEPSIA.**

401—R Bismuthi subcarb., . . . . . ℥ij.  
Morphiæ sulph., . . . . . gr. i-ij.  
Pulv. aromat., . . . . . ℥j. —M.

In chart. no. xii div.

Sig.: A powder in milk before each meal. (*In irritative form.*) BARTHOLOW.

402—R Sodii sulpho-carbolat., . . . . . ℥iv.  
Glycerinæ, . . . . . ℥ij.  
Infusi quassia, . . . . . ℥vj.—M.

Sig.: A tablespoonful before meals. (*In flatulent dyspepsia.*) DELAFIELD.