

DYSPEPSIA (Continued).

403—R Sol. iodinii trichlor. (1-1500), ℥iv.

Sig.: A teaspoonful every two hours. (*When due to presence of bacteria.*)

LANGENBUCH, *Annual Univ. Med. Sci.*

404—R Naphthalini, ℥iss.

Dispensa in capsulas no. xviii.

Sig.: One or two capsules every three to six hours. (*In flatulent dyspepsia and intestinal indigestion.*)

HOLT, *Annual Univ. Med. Sci.*

405—R Ammonii salicylat., ℥ij.
Syr. aurantii cort., ℥j.
Aquæ menthæ pip., ad ℥iv.—M.

Sig.: A tablespoonful half an hour before meals. (*In fermentative dyspepsia.*)

SULLIVAN, *Annual Univ. Med. Sci.*

406—R Bismuthi subnitrat., ℥iv.
Mucil. acaciæ, ℥j.
Sodii bicarb., ℥iv.
Infusi calumbæ, ad ℥viiij.—M.

Sig.: Two tablespoonfuls before meals. (*Irritative dyspepsia, with malnutrition and raw tongue.*)

FOTHERGILL.

407—R Tinct. capsici, ℥xvj.
Tinct. nucis vom., ℥ij.
Tinct. gentianæ co., ad ℥ij.—M.

Sig.: A teaspoonful in water three times daily, with gr. $\frac{1}{5}$ aloin at bed-time, avoiding starchy diet. (*For aggravated dyspepsia with constipation.*)

DA COSTA.

408—R Bismuthi subnitrat.,
Sodii bicarb.,
Pulv. cubebæ, āā ℥j.
Pulv. zingiberis, ℥j.—M.

In pulv. no. xii div.

Sig.: A powder in a wineglassful of water before each meal.

ALONZO CLARK.

409—R Tinct. opii deodorat., gtt. xij.
Magnesii calcinat., gr. xii-xxiv.
Sacchari albi, ℥j.
Aquæ anisi, ℥iss.—M.

Sig.: Shake bottle. One teaspoonful every two hours to a child one year old, until relieved. If much pain, add a little chloroform or Hoffman's Anodyne to the mixture.

J. LEWIS SMITH.