

EPISTAXIS.

432—R Olei erigerontis (canad.), . . . ʒij.

Sig.: Five to fifteen drops on sugar every four hours, or repeated as required.

DE FORREST WILLARD, *Annual Univ. Med. Sci.*

433—R Antipyrin., ʒij.

In capsulas no. xxiv div.

Sig.: One, two, or three to be taken as required. To be used with local treatment.

BEVERLY ROBINSON, *Annual Univ. Med. Sci.*

434—R Pulv. acidi tannici, ʒij.

Sig.: To be insufflated after a small quantity of cocaine has been applied.

FLETCHER INGALS, *Annual Univ. Med. Sci.*

435—R Ext. geranii mac. fld., ʒj.
Aquæ, ʒiij.—M.

Sig.: Syringe the nostrils, or plug with cotton saturated with the fluid.

J. V. SHOEMAKER, *Annual Univ. Med. Sci.*

436—R Ext. hamamelis fld., ʒij.

Sig.: A teaspoonful every one to three hours. If pulse is rapid and bounding, add veratrum viride and morphia. J. V. SHOEMAKER, *Annual Univ. Med. Sci.*

437—R Pulv. ipecac., gr. xx.
Olei theobromæ, ʒss.—M.

Ft. suppositor. no. i.

Sig.: Introduce into the rectum, and when vomiting ceases give—

438—R Pulv. ipecac.,
Ext. glycyrrhizæ, āā ʒss.—M.

Sig.: A powder every three hours.

PEPPER, *Annual Univ. Med. Sci.*

439—R Pulv. aluminis,
Pulv. acidi tannici, āā p. æq.—M.

Sig.: To be insufflated into the nares anteriorly and posteriorly.

SAJOUS.