

ERYSIPELAS.

440—R Acetanilid (antifebrine), . ʒj.

Dispensa in capsulas no. xv.

Sig.: Two capsules as required. (*To reduce the temperature.*)
OSLER, Annual Univ. Med. Sci.

441—R Cretæ præparat.,
Adipis, āā ʒj.
Acidi carbolicæ, ʒj.—M.

Ft. ungt.

Sig.: Apply externally and cover with lint.
SIR DYCE DUCKWORTH, Annual Univ. Med. Sci.

442—R Ichthyol., ʒj.
Vaselini, ʒix.—M.

Sig.: Apply locally.
NUSSBAUM, Annual Univ. Med. Sci.

443—R Ichthyol., ʒss.
Ætheris,
Glycerinæ, āā ʒij.—M.

Sig.: Apply locally.
LORENZ, Annual Univ. Med. Sci.

444—R Sol. hydrarg. bichlor. (3-1000), q. s.

Sig.: Bathe the parts and irrigate the wounds, and cover with iodoform gauze wet with the solution. Apply tar to the red portions of the skin and a little beyond. Then cover with a wet dressing made with "Burow's Fluid."

445—R Aluminis crudi, ʒj.
Plumbi acetat. cryst., ʒv.
Aquæ destillat., ʒxiiss.—M.

Sig.: "Burow's Fluid."
FRAIPONT, Annual Univ. Med. Sci.

446—R Plumbi acetat., ʒj.
Tinct. opii, ʒj.
Aquæ, ad Oj.—M.

Sig.: Shake the bottle well, and wet cloths or lint thoroughly with the lotion, and apply to affected parts.
Charity Hospital, N. Y.

447—R Tinct. ferri chloridi,
Syr. simplicis, āā ʒj.
Aquæ, ʒij.—M.

Sig.: A teaspoonful every two or three hours, well diluted.
Charity Hospital, N. Y.