

**FEVERS, ERUPTIVE AND SIMPLE (Continued).**

455—R Ammonii salicylat., . . . . . ℥j.  
 Syr. simplicis, . . . . . ℥ss.  
 Aquæ menthæ pip., . . . . . ad ℥iiss.—M.

Sig.: A teaspoonful every four hours to a child three years old. (*May be combined with aromatic spirits of ammonia if heart's action is weak.*) To adults, two or three teaspoonfuls may be given every four hours.

SULLIVAN, *Annual Univ. Med. Sci.*

456—R Cocaini hydrochlorat., . . . . . gr. iv.  
 Aquæ destillatæ, . . . . . ℥ij. —M.

Sig.: Seven to fifteen minims hypodermically every two hours. (*In low fevers with nervous depression and weak circulation.*)

DA COSTA, *Annual Univ. Med. Sci.*

457—R Antimon. et potass. tart., . . . . . gr. viij.  
 Aquæ destillatæ, . . . . . ℥viiij.—M.

Sig.: Sponge the scalp frequently. (*For loss of hair after fever.*)

POULAIN, *Annual Univ. Med. Sci.*

458—R Amyl hydratis, . . . . . ℥iij.  
 Syr. simplicis, . . . . . f℥j.  
 Aquæ, . . . . . ad f℥iv.—M.

Sig.: Two tablespoonfuls at bed-time, in water, for adult. (*For insomnia of fevers.*)

VON MERING, *Annual Univ. Med. Sci.*

459—R Antifebrin., . . . . . ℥i-ij.  
 Spts. vini gallici, . . . . . ℥ss.  
 Syrupi simplicis, . . . . . ad f℥ij.—M.

Sig.: A teaspoonful every four hours, or as required. (*To reduce the temperature.*)

HEINZELMANN, *Annual Univ. Med. Sci.*

460—R Antipyrin., . . . . . ℥j.  
 Syr. simplicis, . . . . . ℥ss.  
 Aquæ cinnamomi, . . . . . ad ℥ij.—M.

Sig.: One-half to one teaspoonful every hour or two for children.

PENZOLDT.

461—R Sodii bromidi, . . . . . gr. x-xx.  
 Syr. aurantii cort., . . . . . ℥ss.  
 Aquæ, . . . . . ad ℥iiss.—M.

Sig.: A teaspoonful every quarter of an hour for children.

A. A. SMITH.