

FEVERS, ERUPTIVE AND SIMPLE (Continued).

462—R Liq. ammonii acetat., ℥iiiiss.
Spts. ætheris nitrosi, ad ℥iv.—M.

Sig.: A teaspoonful to a tablespoonful, according to age. HARTSHORNE.

463—R Acetphenetid., ℥i—iss.
Sacchari alb., ℥ss. —M.

In pulv. no. xx div.

Sig.: One every four to six hours, as required, for fever in children. (*For adults twice the dose.*)

KOBLER, Annual Univ. Med. Sci.

FEVER, HECTIC.

464—R Tinct. digitalis, ℥iij.
Tinct. ferri chlor., ℥v.—M.

Sig.: Fifteen drops three or four times daily, well diluted. BARTHOLOW.

465—R Quiniæ sulph., ℥j.

In pulv. no. xii div.

Sig.: A powder, in water, three or four times daily. PHILLIPS.

466—R Syr. calcis lactophosphat., ℥iv.

Sig.: A teaspoonful three or four times daily.

BENEKÉ.

467—R Antipyrin., ℥ij.
Aquæ, ℥viiij.—M.

Sig.: Two tablespoonfuls, followed by one tablespoonful every hour till temperature is normal. Continue for four or five days. If the solution is not strong enough, double it. On the sixth or seventh day omit the second dose. If three doses have been given daily, omit the third. After a few days longer, drop the second.

PRIBRAM, Annual Univ. Med. Sci.

FEVERS, INTERMITTENT AND REMITTENT.

468—R Quiniæ muriatis, gr. vj.
Aquæ bullientis, ℥xij.—M.

Sig.: Inject deeply into the tissues four to six minims of the hot solution.

PULAWSKI, Annual Univ. Med. Sci.