

FEVERS, INTERMITTENT AND REMITTENT (Continued).

469—R Acidi arseniosi, gr. j.
Extracti taraxaci, ℥j.—M.

Ft. massa et in pil. no. xl div.

Sig.: One to four pills every three or four hours until twenty or thirty pills are taken. (*In hemorrhagic malarial fever.*)

RIGGS, Annual Univ. Med. Sci.

470—R Pulv. opii, gr. xij.
Pulv. capsici, gr. xxxvj.
Quiniæ sulph., ℥j. —M.

In pulveres no. xii div.

Sig.: One powder three times daily. (*In inveterate forms.*)

ALONZO CLARK.

471—R Ext. ergotæ aq. (Bonjean), . . gr. xxx.
Aquæ destillatæ,
Glycerinæ, āā ℥iiss.—M.

Sig.: Inject an hypodermic syringeful into the spleen once or twice, at intervals of a few days.

ROUQUETTE, Annual Univ. Med. Sci.

472—R Olei phosphorati, ℥ss.

Sig.: Five drops, well diluted, three times daily.

SOZINSKEY, Annual Univ. Med. Sci.

473—R Ammonii picratis, ℥j.

Ft. massa et in pilulas no. xl div.

Sig.: One to three pills four or five times daily.

H. M. CLARK, Annual Univ. Med. Sci.

474—R Potassii permanganatis, . . . gr. xij.
Aquæ destillatæ, ℥iij. —M.

Sig.: One or two teaspoonfuls three times daily. (*In chronic malarial cases.*)

JOS. LEVI, Annual Univ. Med. Sci.

475—R Antipyrin., ℥iv.
Syr. toltutan., ℥iiss.
Aquæ, ad ℥iij.—M.

Sig.: One or two teaspoonfuls thrice daily.

ANTONY, Annual Univ. Med. Sci.