

FEVERS, INTERMITTENT AND REMITTENT (Continued).

476—R Acidi carbolici, gr. xlvij.
Syr. acaciæ, ℥iiss.
Aquæ cinnamomi, ad ℥iv.—M.

Sig.: A teaspoonful every three hours until five doses are taken daily.

LUZZATTO, *Annual Univ. Med. Sci.*

477—R Tinct. iodinii, ℥vj.
Potassii iodidi, ℥ij.
Aquæ cinnamomi, ad ℥iij.—M.

Sig.: A teaspoonful three times daily.

Charity Hospital, N. Y.

478—R Acidi carbolici, ℥j.
Tinct. iodinii co., ℥iij.—M.

Sig.: Four drops every four hours, well diluted.

BARTHOLOW.

479—R Quiniæ sulphat., ℥iv.
Acidi sulphurici dil., q. s. ut ft. sol.
Spts. ætheris nitrosi, ℥ss.
Syr. tolutani,
Aquæ, āā q. s. ad ℥ij.—M.

Sig.: A teaspoonful three or four times daily.

DA COSTA.

480—R Quiniæ sulphat., gr. xvj.
Ext. glycyrrhizæ, ℥j.
Syr. rubi idæi (raspberry), ℥ij.—M.

Sig.: A teaspoonful three to five times daily for a child three years old.

J. LEWIS SMITH.

481—R Chinoidini, ℥ij.
Resinæ podophylli, gr. iv.
Ferri sulph. exsic., ℥j.—M.

Ft. massa et in pil. no. xx div.

Sig.: One three times daily.

BARTHOLOW.

FEVER, TYPHOID.

482—R Hydrarg. chlorid. mit., gr. xxij.
In pulv. no. iii div.

Sig.: One powder every half-hour till three are taken. (*Only during the first week.*)

VON ZIEMSEN, *Annual Univ. Med. Sci.*