

FEVER, TYPHOID (Continued).

483—R Acidi sclerotici, ℥j.
Aquæ destillatæ, ℥v.—M.

Sig.: A hypodermic syringe-ful every half-hour as a
styptic. (*In typhoid intestinal hemorrhage.*)
VON ZIEMSEN, *Annual Univ. Med. Sci.*

484—R Bismuthi subnitratæ,
Pepsinæ, āā ℥ij.—M.

In pulv. no. xii div.
Sig.: Four powders daily, at suitable intervals, in
milk. ROOSEVELT HOSPITAL, *Annual Univ. Med. Sci.*

485—R Antifebrin., ℥ss-j.
Elixir simplicis, ℥j.—M.

Sig.: A teaspoonful when required. (*For pyrexia.*)
BEREZOVSKY, *Annual Univ. Med. Sci.*

486—R Tinct. strophanth. hisp. (1-20), ℥ss.

Sig.: Four to eight minims three or four times
daily. (*For weak heart of typhoid.*)
QUINLAN, *Annual Univ. Med. Sci.*

487—R Thallin sulphat., gr. xxxij.
Aquæ menthæ pip., ℥j.—M.

Sig.: A teaspoonful when required. (*For pyrexia
of typhoid.*)
MINOT, *Annual Univ. Med. Sci.*

488—R Thymoli, ℥j.
Ext. gentianæ, q. s.

Ft. massa et in pil. no. xxiv div.
Sig.: Two pills every six hours. (*For diarrhœa
and dry tongue of typhoid.*)
HENRY, *Annual Univ. Med. Sci.*

489—R Antipyrin., ℥viii.
Syr. tolutani, ℥j.
Aquæ, ad ℥ij.—M.

Sig.: Two or three teaspoonfuls at first, and one
teaspoonful hourly thereafter, until temperature is
reduced. (*For pyrexia of typhoid.*)
MINOT, *Annual Univ. Med. Sci.*

490—R Pulv. pepsinæ cryst., ℥ij.
Acidi muriatici dil., ℥ss.
Syr. simplicis, ℥j.
Aquæ, q. s. ad ℥ij.—M.

Sig.: A teaspoonful after food three times daily.
(*For antifebrile action, and for the impaired digestion
of typhoid.*)
WITHERSTINE.