

FEVER, TYPHUS.

491—R Quiniæ sulphat., ℥iv.
Acidi sulphurici dil., q. s. ut ft. sol.
Syr. simplicis, ℥ss.
Aquæ, ad ℥j.—M.

Sig.: A teaspoonful every two hours until fever is lessened. GOOLDEN.

492—R Acidi phosphorici dil., ℥j.

Sig.: Twenty to thirty minims, well diluted, every four or six hours. TANNER.

493—R Ext. opii, gr. vj.

In pil. no. xii div.

Sig.: One pill every three hours, until quiet. (*For restlessness, when there is no lung complication present.*) TANNER.

494—R Moschi, gr. x.
Ætheris,
Tinct. opii, āā ℥xx.
Aquæ cinnamomi, ℥j. —M.

Fiat haustus.

To be taken at one draught. (*In last stages of typhus.*) HOOPER.

495—R Tinct. belladonnæ, ℥ss.
Tinct. aconiti rad., ℥ss.—M.

Sig.: Ten drops every two hours. (*For rapid pulse and dry tongue.*) HARLEY.

496—R Tinct. opii, ℥j.
Antimon. et potass. tart., gr. iv.
Aquæ camphoræ, ℥viiij.—M.

Sig.: A tablespoonful every two hours. (*For restlessness and insomnia.*) GRAVES.

FEVER, YELLOW.

497—R Pilocarpinæ muriat., gr. iij.
Aquæ destillatæ, ℥ij. —M.

Sig.: Ten minims hypodermically. (*To favor diaphoresis.*) HEBERSMITH, Annual Univ. Med. Sci.