

GASTRALGIA (See Neuralgia, Catarrh, Colic, Dyspepsia).

GASTRIC ULCER (See Ulcer).

GLANDS, ENLARGED LYMPHATIC.

538—℞ Syr. ferri iodidi, ℥j.
Sig.: Five to forty minims, according to age, well
diluted, after meals, internally. BARTHOLOW.

539—℞ Potassii iodidi, ℥i-iv.
Syr. aurantii cort., ℥j.
Aquæ cinnamomi, ad ℥ij.—M.
Sig.: A teaspoonful in water three times daily.
RINGER.

540—℞ Calcii sulphidi, gr. vj.
In pil. no. xxiv div.
Sig.: One pill every four to six hours. RINGER.

541—℞ Hydrarg. cum cretæ, gr. iv.
Sacch. lactis, ℥j. —M.
In pulv. no. xx div.
Sig.: One every two hours. (*In acute stage, ton-*
sillitis, parotiditis, etc.) BARTHOLOW.

542—℞ Tinct. iodinii, ℥ss.
Sig.: Apply with a brush to the part.
T. M. MARKOE.

543—℞ Ungt. plumbi iodidi, ℥j.
Sig.: Apply locally. BARTHOLOW.

544—℞ Ungt. iodinii co., ℥j.
Sig.: Apply locally. H. B. SANDS.

GLEET (See Gonorrhœa).

GOITRE (See also Glands, Enlargement of).

545—℞ Ungt. hydrarg. iodidi rubr., ℥j.
Sig.: Rub in a piece the size of a pea, and expose to
sun's rays. RINGER.

546—℞ Tinct. iodinii co., ℥j.
Sig.: Apply locally with a brush. Also five to fifteen
minims in water three times daily, internally.
BARTHOLOW.