

GOUT (Continued).

569—R Paraldehyde, ℥ss.
Syr. simplicis, ℥iiss.—M.

Sig.: A teaspoonful to a tablespoonful, well diluted,
when required. (*For gouty insomnia.*)

G. F. HODGSON.

570—R Colchicini, gr. j.
Ext. colocynth. co., ℥ss.
Quiniæ sulph., ℥iij.—M.

Ft. massa et in pil. no. lx div.

Sig.: One pill every four hours.

BARTHOLOW.

571—R Ext. colchici acet. (B. P.),
Ext. opii aquosi, āā gr. xv.
Potassii iodidi, ℥iv.
Potassii acetatis, ℥ij.
Aquæ destillatæ, ℥iiss.
Vini albi, ℥ss.—M.

Sig.: Twenty drops thrice daily.

LALLEMAND.

572—R Potassii carbonatis,
Potassii nitratis, āā ℥iiss.
Aquæ, ℥viiij.—M.

Sig.: A tablespoonful thrice daily. (*In gouty attacks.*)

573—R Vini colchici rad., ℥j.
Magnesiæ calcin. (Husband's), ℥j.
Aquæ menthæ pip., ℥iv.—M.

Sig.: A tablespoonful thrice daily. (*In gouty rheumatism.*)

574—R Tinct. rhei et sennæ, ℥iiss.
Syr. zingiberis, ℥iij.
Tinct. opii, ℥j.—M.

Sig.: A teaspoonful at once, in hot water. (*In spasmodic gout.*)

HARTSHORNE.

575—R Magnesii sulph., ℥ij.
Potassii bicarb., gr. xv.
Tinct. colchici sem., ℥x.
Infusi buchu, ℥j.—M.

Ft. haustus.

Sig.: To be taken every four or six hours, followed
by a large draught of water, not too cold. (*In acute gout.*)

FOTHERGILL.