

HEMORRHOIDS (Continued).

658—R Tinct. hamamelis, . . . . . ℥iv.

Sig.: Inject into the rectum one-half to one tea-  
spoonful in an ounce of cold water daily before rising.  
Also take internally two to five minims three times  
daily. RINGER.

659—R Potassii bromidi, . . . . . ℥iij.  
Glycerinæ, . . . . . ℥iiss.—M.

Sig.: Apply locally to ease pain. RINGER.

660—R Acidi nitrici, . . . . . ℥ss-j.  
Aquæ, . . . . . ℥viiij.—M.

Ft. lotio.

Sig.: Apply as a wash. (*In bleeding piles.*)  
RINGER.

661—R Tinct. nucis vomicæ, . . . . . ℥j.  
Ext. ergotæ fld., . . . . . ℥j.

Sig.: A teaspoonful three or four times daily. (*For  
bleeding piles and post-partum piles.*) BARTHOLOW.

662—R Ext. belladonnæ, . . . . . ℥j.  
Cerati simplicis, . . . . . ℥j.—M.

Ft. ungt.

Sig.: Use as an ointment. (*For painful piles.*)  
HARTSHORNE.

663—R Acidi tannici, . . . . . gr. xx-xxx.  
Aquæ, . . . . . ℥vj. —M.

Sig.: To be injected, after being cooled with ice, into  
the rectum. (*For bleeding piles.*) HARTSHORNE.

664—R Ext. opii, . . . . . gr. viij.  
Ext. belladonnæ, . . . . . gr. iv.  
Olei theobromæ, . . . . . ℥iv.—M.

Ft. massa et in suppositoria no. viii div.

Sig.: One to be introduced into the bowel every  
three or four hours. (*For painful piles.*)

WITHERSTINE.

HEPATITIS (See Catarrh and Biliousness).

HERPES (See Skin Diseases).