

INCONTINENCE OF URINE (Continued).

703—R Acidi benzoici, ℥ij.
Aquæ cinnamomi, ℥vj.—M.
Sig.: A tablespoonful thrice daily. HARTSHORNE.

704—R Tinct. ferri muriatis, ℥j.
Decocti uvæ ursæ, ℥vj.—M.
Sig.: A tablespoonful two or three times daily.
WILLIS.

705—R Tinct. belladonnæ, ℥j.
Sig.: Ten to twenty drops thrice daily. RINGER.

706—R Santonini, gr. xvj.
Olei ricini, ℥j. —M.
Sig.: One or two teaspoonfuls before breakfast, for
two or three mornings. RINGER.

707—R Collodii, ℥ss.
Sig.: Put a drop on the meatus to seal it, at bed-
time. Remove with finger-nail in morning.
D. CORRIGAN.

708—R Strychniæ, gr. j.
Acidi acetici, gtt. ij.
Sacchari albi, ℥ij.
Aquæ destillatæ, ℥ij.—M.
Ft. solutio.
Sig.: Fifteen to thirty drops for a child six to twelve
years old. MAGENDIE.

INFLUENZA (See Catarrh and Hay Fever).

INGROWING TOE-NAIL.

709—R Acidi tannici, ℥j.
Aquæ destillatæ, ℥vj.—M.
Sig.: Paint soft parts twice daily.
MIALI, Annual Univ. Med. Sci.

710—R Pulv. plumbi acetat., ℥j.
Tinct. opii, ℥ij.
Aquæ, ad ℥viiij.—M.
Sig.: Shake well, and apply constantly until the
inflammation is reduced and pain alleviated; then
separate the granulating surface from the nail and
insert a small pledget of cotton; then use—