

LARYNGITIS.

761—R Tinct. aconiti radice, . . . . . ℥ss.

Sig.: One drop every hour in water. Best results when following a dose of castor oil. When it has existed some days, then give—

762—R Vini Mariani, . . . . . Oj.

Sig.: A wineglassful every three hours, with absolute rest of voice. (*In acute laryngitis.*) SAJOUS.

763—R Acidi benzoici, . . . . . gr. ss.  
Sodii biboratis, . . . . . gr. iss.  
Acaciæ, sugar, or currant-paste, q. s. —M.

Ft. trochiscum no. i.

Sig.: One every hour. (*In acute laryngitis.*) SAJOUS.

764—R Sodii biboratis, . . . . . gr. viij.  
Aquæ, . . . . . ℥ij.  
Aquæ cologniensis, . . . . . gtt. x.—M.

Sig.: Use frequently as a spray with atomizer. (*Chronic form.*) SAJOUS.

765—R Potassii permanganatis, . . . . . gr. ij.  
Aquæ destillatæ, . . . . . ℥ij. —M.

Sig.: Use with atomizer several times daily. (*In fetid variety of chronic laryngitis.*) SAJOUS.

766—R Argenti nitratis, . . . . . gr. lx.  
Aquæ destillatæ, . . . . . ℥j. —M.

Sig.: Apply locally on cotton after using—

767—R Sol. cocaini muriatis (10 per cent.), . . . . . ℥j.

Sig.: Apply locally to the larynx. (*In chronic form.*) SEILER.

768—R Sol. cocaini muriatis (15–20 per cent.), . . . . . ℥ss.

Sig.: Apply locally to the larynx before and, if much pain, after the following:—

769—R Acidi lactici, . . . . . ℥ss.

Sig.: Apply locally to all infiltrations and ulcerations in the larynx. (*In tuberculous laryngitis.*)

HERING, Annual Univ. Med. Sci.