

LOCOMOTOR ATAXIA (Continued).

800—R Antipyrin., ℥ij.
Syr. zingiberis, ℥ij.
Aquæ cinnamomi, ad ℥iv.—M.

Sig.: A teaspoonful every one to four hours for three to six doses. (*In lightning-pain of locomotor ataxia.*)
GERMAIN SÉE, Annual Univ. Med. Sci.

801—R Antifebrin., ℥j.

Dispensa in capsulas no. xv.

Sig.: One or two capsules every half-hour for two doses, if necessary; then one every four or six hours if required. (*For pains of locomotor ataxia.*)

DUJARDIN-BEAUMETZ, Annual Univ. Med. Sci.

LUMBAGO.

802—R Pulv. potassii nitratis, ℥ij.

In pulv. no. xii div.

Sig.: A powder in a half-tumblerful of water every hour or two. (*When urine is scanty and high colored.*)

RINGER.

803—R Olei terebinthinæ, ℥ii-ij.
Mucilag. acaciæ, q. s. ut ft. emuls.
Syr. zingiberis, ℥j.
Aquæ, ad ℥iij.—M.

Sig.: A tablespoonful every four to six hours, carefully, lest strangury and nephritis supervene. (*When urine is clear and abundant, and bowels regular.*)

WARING.

804—R Atropiæ sulphatis, gr. i½.
Morphiæ sulphatis, gr. xvj.
Aquæ destillatæ, ℥j. —M.

Sig.: Five minims injected deep into the muscular tissues.

DA COSTA.

805—R Antipyrin., ℥ij.
Syr. tolutani, ℥ij.
Aquæ menthæ pip., ad ℥iv.—M.

Sig.: A teaspoonful every one to four hours for three to six doses.

GERMAIN SÉE, Annual Univ. Med. Sci.

806—R Methyl chloridi, ℥ss.

Sig.: Use locally, applying carefully.

DEBOVE, Annual Univ. Med. Sci.