

**LUMBAGO (Continued).**

807—R Tinct. actææ racemos. (1 to 4), ℥ss.

Sig.: Five minims every hour, or fifteen to thirty minims thrice daily. RINGER.

808—R Aquæ destillatæ, . . . . . ℥j.

Sig.: Thirty to sixty minims hypodermically. BARTHOLOW.

809—R Empl. belladonnæ (6 in. x 4 in.).

Sig.: Apply locally. (*For persistent remains, affecting a small spot.*) RINGER.

810—R Potassii iodidi, . . . . . ℥iv.  
Vini colchici sem., . . . . . ℥j.  
Syr. sarsaparillæ co., . . . . . ℥j.  
Aquæ, . . . . . q. s. ad ℥iij.—M.

Sig.: A teaspoonful in a wineglassful of water every four hours. J. M. LEEDOM.

811—R Tinct. iodinii, . . . . . ℥ij.  
Tinct. aconiti rad., . . . . . ℥iij.  
Chloroformi, . . . . . ℥iv.  
Linimenti saponis co., . . . . . ad ℥iij.—M.

Sig.: Apply every few hours locally. *Bellevue Hosp., N. Y.*

**LUPUS.**

812—R Acidi chromici, . . . . . gr. c.  
Aquæ destillatæ, . . . . . ℥j. —M.

Sig.: Apply locally. BARTHOLOW.

813—R Zinci sulphatis exsiccati, . . . . . ℥ij.

Sig.: Dust on the surface, and, when it is cauterized, remove the slough with a poultice. BARTHOLOW.

814—R Zinci chloridi,  
Antimonii chloridi, . . . . . āā ℥j.  
Pulv. iridis florentinæ rad., . . . . . gr. x.  
Acidi hydrochlorici puri, . . . . . ℥x.—M.

Sig.: Use as a caustic, spread on linen strips a quarter-inch wide. Lay the strips on the spot, cover with lint, and leave for twenty-four hours.

*KAPOSI, Annual Univ. Med. Sci.*