

**MENINGITIS (Continued).**

860—R Hydrarg. chloridi mitis,  
Pulv. jalapæ,  
Sacchari albi, . . . . . āā ʒj.—M.

In pulv. no. x div.

Sig.: A powder every hour until free purgation occurs. (*In cerebro-spinal meningitis.*)

ROBERT, Annual Univ. Med. Sci.

861—R Morphiæ sulphatis, . . . . . gr. ij.  
Aquæ destillatæ, . . . . . ʒj. —M.

Sig.: Five minims hypodermically every three to five hours, or more often, to control the vomiting and to keep the patient quiet and comfortable. Absolute rest, and small quantities of strong food frequently per os and per rectum. (*In cerebro-spinal form.*)

LEYDEN, Annual Univ. Med. Sci.

862—R Sodii bromidi, . . . . . ʒij.  
Chloral hydratis, . . . . . ʒj.  
Syr. aurantii cort., . . . . . ʒj.  
Aquæ, . . . . . q. s. ad ʒiij.—M.

Sig.: A dessertspoonful every hour or two until excitement abates. HERRMANN, Annual Univ. Med. Sci.

863—R Morphiæ sulphatis, . . . . . gr. ss.  
Acidi sulphurici aromat., . . . . . ʒj.  
Elixir cinchonæ, . . . . . ad ʒvj.—M.

Sig.: A teaspoonful every two hours for a child twelve years old. (*Cerebro-spinal form.*)

MEIGS and PEPPER.

864—R Tinct. aconiti radice, . . . . . ʒij.  
Tinct. opii deodoratæ, . . . . . ʒv.—M.

Sig.: Seven drops in water, every two hours, during the stage of excitement. (*Cerebral meningitis.*)

BARTHOLOW.

865—R Acidi hydrocyanici dil., . . . . . ℥xx-xl.  
Sodii bicarbonatis, . . . . . ʒiii-v.  
Syr. simplicis,  
Aquæ, . . . . . q. s. ad ʒiiss.—M.

Sig.: A teaspoonful every three or four hours for severe vomiting. (*Cerebro-spinal form.*)

DELAFIELD.

866—R Tinct. ferri chloridi, . . . . . ʒij.

Sig.: Twenty to thirty minims every two hours.

KLAPP.

**MENINGITIS, CEREBRO-SPINAL (See Meningitis).**