

**MENORRHAGIA.**

867—R Ext. ipecac. fld., . . . ʒij.  
Ext. ergotæ fld., . . . ʒiv.  
Ext. digitalis fld., . . . ʒij.—M.

Sig.: A half to one teaspoonful as required, until emesis occurs. BARTHOLOW.

868—R Ext. gossypii fld.,  
Syr. simplicis, . . . āā ʒj.—M.

Sig.: A teaspoonful every four hours. PARVIN.

869—R Olei erigerontis, . . . ʒj.

Sig.: Five drops on a lump of sugar every three or four hours. BARTHOLOW, Annual Univ. Med. Sci.

870—R Ext. geranii maculat. fld., . ʒiv.

Sig.: A teaspoonful every hour for a few doses, then every three to four hours. May be used with advantage locally. SHOEMAKER, Annual Univ. Med. Sci.

871—R Ext. rhois aromat. fld., . . ʒj.

Sig.: Fifteen to sixty minims thrice daily. UNNA, Annual Univ. Med. Sci.

872—R Ext. hydrastis can. fld., . . ʒj.

Sig.: Twenty drops four times daily. R. W. WILCOX, Annual Univ. Med. Sci.

873—R Acidi gallici, . . . ʒss.  
Acidi sulphurici dil., . . . ʒj.  
Tinct. opii deod., . . . ʒj.  
Inf. rosæ comp., . . . ʒiv.—M.

Sig.: A tablespoonful every four hours or oftener. BARTHOLOW.

874—R Tinct. hamamelis, . . . ʒij.

Sig.: One-half to one teaspoonful thrice daily. RINGER.

875—R Pulv. potassii bromidi, . . ʒij.

In pulv. no. xii div.

Sig.: A powder in a wineglassful of water three times daily. (*In flooding of young women who menstruate too often, as well as too copiously.*) Begin before the period and continue till it is over. RINGER.