

**MENORRHAGIA (Continued).**

876—R Ext. ergotæ fld., . . . . . ℥ij.  
Sig.: A half to one teaspoonful thrice daily.

877—R Liq. ferri perchloridi, . . . . . ℥iv.  
Aquaë, . . . . . ℥xij.—M.  
Sig.: Inject slowly and carefully into the uterus with a Davidson's syringe fitted with a long uterine tube. Avoid introducing air. Allow a free outlet for the fluid. (*In post-partum hemorrhage.*) R. BARNES.

878—R Tinct. sabinæ, . . . . . ℥ss.  
Sig.: Five to ten drops in cold water every half to three hours. PHILLIPS.

**MERCURIALISM (See Ptyalism).**

**METRITIS.**

879—R Tinct. aconiti rad., . . . . . gtt. xvj.  
Ext. gelsemii fld., . . . . . ℥j.  
Ext. ergotæ fld., . . . . . ad ℥j.—M.  
Sig.: A teaspoonful every two to six hours. (*Also in uterine tumor.*) BARTHOLOW.

880—R Tinct. iodinii co., . . . . . ℥j.  
Sig.: Use locally on a probe wrapped with absorbent cotton, once or twice weekly. Two applications are made, and a glycerin tampon is left against the cervix. In the intervals, let the patient use a gallon or more of hot water as a vaginal injection twice or thrice daily. T. G. THOMAS.

**MIGRAINE (See Headache and Neuralgia).**

**MITRAL DISEASE (See Heart-Disease).**

**MORNING SICKNESS (See also Vomiting).**

881—R Tinct. cantharidis,  
Tinct. ferri muriatis, . . . . . āā f℥j.—M.  
Sig.: Twenty-five drops, well diluted, three times daily. HIGGINS, Annual Univ. Med. Sci.

882—R Sodii bicarbonatis, . . . . . gr. iv.  
Acidi hydrocyanici dil., . . . . . gtt. j.  
Syr. lactopeptini, . . . . . ℥j. —M.  
Sig.: To be given half an hour before meals. J. FREE, Annual Univ. Med. Sci.