

**OPHTHALMIA (Continued).**

941—R Hydrarg. chloridi corrosiv., . gr. j.  
Aquaë, . . . . . ℥xij.—M.

Sig.: Irrigate the eye frequently with the solution.  
CREDÉ.

942—R Acidi boracici, . . . . . gr. xvj.  
Acidi salicylici, . . . . . gr. ij.  
Glycerinæ, . . . . . ℥xl.  
Aquaë bullientis, . . . . . ad ℥j. —M.

Sig.: Instill into the eye, after cauterizing trachoma follicles with the thermo-cautery. (*In trachoma.*)  
ARMAIGNAC, Annual Univ. Med. Sci.

943—R Pulv. acidi tannici, . . . . . ℥j.  
Pulv. acidi boracici, . . . . . ℥ij.—M.

Sig.: Dust the lids. Cautery may precede it. (*In trachoma.*)

WICHERKIEWICZ, Annual Univ. Med. Sci.

944—R Hydrarg. chloridi corrosiv., . gr. j.  
Aquaë, . . . . . ℥j.—M.

Sig.: Apply once daily to the lids with a brush. Use (1-7000) sol. several times daily.

QUAITA, Annual Univ. Med. Sci.

945—R Cocaini sulphatis, . . . . . gr. iv.  
Atropiæ sulphatis, . . . . . gr. ss.  
Vaselini, . . . . . ℥v.—M.

Sig.: To be applied with a camel's hair brush. The chemosis and pain are relieved instantly. (*Catarrhal ophthalmia.*)

LEAHY.

946—R Sodii biboratis, . . . . . gr. xx.  
Aquaë camphoræ, . . . . . ℥iss.  
Mucil. cydonii sem., . . . . .  
Aquaë lauro-cerasi, . . . . . āā ℥ss.—M.

Sig.: Apply to the lids and drop into the eyes three times daily. (*Catarrhal blepharitis.*)

947—R Hydrarg. oxidi rubri, . . . . . gr. vj.  
Plumbi subacetat. cryst., . . . . . gr. ij.  
Vaselini, . . . . . ℥v. —M.

Ft. ungt.

Sig.: Apply to the free border of the eyelids once daily, after bathing the eyelids in hot water. (*In chronic blepharitis.*)

PARINAUD.