

**PARALYSIS (Continued).**

992—℞ Coniæ, . . . . . ℥iii-℥xij.  
Acidi acetici fort., . . . . . ℥iii-℥xij.  
Spts. vini rectificati, . . . . . ℥j.  
Aquæ destillatæ, . . . . . ad ℥ij.—M.

["Add the acid carefully until neutralization, using test-paper. If turbid, add more alcohol."]

Sig.: Begin with one minim, and gradually increase as necessary. Five minims contain one minim of conia. (*In blepharospasm and paralysis agitans.*)

BURMAN.

993—℞ Strychniæ sulphatis, . . . . . gr. ij.  
Aquæ destillatæ, . . . . . ℥j.—M.

Sig.: One to five minims hypodermically. (*In infantile paralysis, diphtheritic and lead paralyses, aphonia, reflex paraplegia, and hysterical paralysis. In chronic hemiplegia and local paralyses, as facial, sphincteric, etc.*)

BARTHOLOW.

994—℞ Ammonii iodidi, . . . . . ℥j.  
Ammonii carbonatis, . . . . . ℥ij.  
Liq. ammonii acetatis, . . . . . ℥vj.—M.

Sig.: A tablespoonful thrice daily. (*To absorb thrombi in incipient hemiplegic paralysis, due to endarteritis deformans.*)

BARTHOLOW.

**PARTURITION (See Labor).**

**PEDICULI (See Lice).**

**PEMPHIGUS (See Skin Diseases).**

**PERICARDITIS (See also Heart-Disease).**

995—℞ Tinct. aconiti radiceis, . . . . . ℥ss.

Sig.: Half a drop to a drop in a teaspoonful of water every ten minutes or quarter of an hour for two hours; then every hour or two. If much prostration, a smaller dose.

RINGER.

996—℞ Hydrarg. chloridi mitis,  
Pulv. ipecac., . . . . . āā gr. vj.  
Potassii nitratis, . . . . . ℥ss-j.—M.

In pulv. no. xii div.

Sig.: A powder every three hours.

HARTSHORNE.