

PERICARDITIS (Continued).

997—R Antimonii et potassii tart., . . . gr. iv.
Tinct. opii, ℥j.
Aquæ camphoræ, ℥viiij.—M.

Sig.: A tablespoonful every two hours. (*In acute form.*) GRAVES.

998—R Empl. cantharidis, 2 in. x 3 in.

Sig.: Apply over the præcordial space. Repeat at intervals after the skin is healed. (*In chronic stage.*) TANNER.

PERIOSTITIS (NODES).

999—R Potassii iodidi,
Potassii carbonatis, āā ℥ij.
Spts. vini rectificati, ℥ij.
Aquæ, ℥xj.—M.

Fiat lotio.

Sig.: Apply twice daily (*in early stage*), with the following internally:—

1000—R Potassii iodidi,
Potassii chloratis, āā ℥j.
Potassii bicarbonatis, ℥iiij.—M.

In chartulas no. xii div.

Sig.: One powder morning and evening in half a pint of milk. ERICHSEN.

1001—R Sodii iodidi, ℥j.
Decocti sarsaparillæ co., ℥viiij.—M.

Sig.: One-sixth part three times daily. TANNER.

1002—R Potassii iodidi, ℥j.
Syr. aurantii cort., ℥j.
Aquæ aurantii flor., ℥v.—M.

Sig.: A tablespoonful twice daily in hop-tea.

LISFRANC.

1003—R Potassii iodidi, ℥ij.
Ammonii iodidi, ℥j.
Tinct. cinchonæ co., ℥iiij.—M.

Sig.: A teaspoonful, largely diluted with water, after eating. VAN BUREN and KEYES.