

PERIOSTITIS (Continued).

1004—R Cadmii iodidi, . . . . . ℥j.  
Adipis preparatæ, . . . . . ℥j.  
Linimenti aconiti, . . . . . ℥ij.—M.  
Ft. ungt. TANNER.

1005—R Zinci iodidi, . . . . . ℥j.  
Adipis, . . . . . ℥j.—M.  
Ft. ungt.  
Sig.: Apply twice daily. HOOPER.

1006—R Morphiæ, . . . . . gr. viij.  
Hydrarg. oleatis (10 per cent.  
ad 20 per cent.), . . . . . ℥j.—M.  
Sig.: Apply with a brush. JOHN MARSHALL.

1007—R Potassii iodidi, . . . . . ℥j.  
Aquæ bullientis, . . . . . ℥j.  
Vaselini, . . . . . ℥vij.—M.  
Ft. ungt.  
Sig.: Apply twice daily, and use the following:—

1008—R Potassii iodidi, . . . . . ℥i-ij.  
In pulv. no. xii div.  
Sig.: One powder, morning and evening, in a glassful  
of milk. RINGER.

1009—R Potassii iodidi, . . . . . gr. ii-x.  
Potassii bromidi, . . . . . gr. v-xx.  
Ammonii carbonatis, . . . . . gr. v.  
Spts. chloroformi, . . . . . ℥xv.  
Aquæ, . . . . . q. s. ad ℥j.—M.  
Sig.: To be taken three times daily. BERKELEY HILL.

1010—R Iodini, . . . . . gr. ss.  
Potassii iodidi, . . . . . ℥ss.  
Syr. papaveris, . . . . . ℥ss.  
Infusi gentianæ co., . . . . . ℥x.—M.  
Sig.: Take two tablespoonfuls thrice daily. Take a  
half-grain of morphia acetate at night. (*In weakly  
constitutions.*) BRANSBY COOPER.