

**PERITONITIS.**

1011—R Morphiae sulphatis, . . . gr. viij.  
Aquaë destillatæ, . . . ℥iv.— M.

Sig.: Begin with a dessertspoonful and wait two hours. If no effect, give three teaspoonfuls and wait two hours. If still no effect, give four teaspoonfuls and wait two hours. The medicine should be increased gradually to produce these effects: to allay pain, to produce gentle sleep, to reduce the respirations to twelve per minute when aroused (may get as low as eight, but should go no lower). Continue these effects for two days, and then gradually diminish the dose, but if the symptoms return, increase again.

ALONZO CLARK.

1012—R Tinct. aconiti rad., . . . ℥ij.  
Tinct. opii deodorat., . . . ℥vj.—M.

Sig.: Eight drops in water every hour or two. Increase the laudanum if the pain is severe.

BARTHOLOW.

1013—R Antimonii et potassii tart., . . gr. j.  
Tinct. opii, . . . ℥j.  
Aquaë camphoræ, . . . ℥viiij.—M.

Sig.: A tablespoonful every two hours. (*When nervous excitement is present.*)

GRAVES.

1014—R Tinct. aconiti folii, . . . ℥v.  
Ext. veratri viridis fld., . . . ℥j.—M.

Sig.: Take twelve drops every two hours. (*Where opium is inadmissible.*)

ELLIS.

1015—R Pulv. piperis,  
Pulv. zingiberis, . . . āā ℥j.  
Sinapis nigræ contus., . . . ℥viiij.  
Aquaë bullientis, . . . q. s.—M.

Ft. cataplasma. (*As a counter-irritant.*) ELLIS.

1016—R Acetphenetidin., . . . ℥ij.

In pulv. no. xii div.

Sig.: A powder stirred in a little water, as required. One-third to one-half the dose for children. (*For febrile condition.*)

KOBLER, *Annual Univ. Med. Sci.*

1017—R Acidi tannici, . . . gr. iij-clxxx.  
Glycerinæ, . . . q. s. ad ft. sol.—M.

Sig.: To be taken in divided doses during the day. (*In localized peritonitis.*)

DEBOUÉ, *Annual Univ. Med. Sci.*